

ORAL SUSPENSION

Calcimax[®] Energy

More Calcium, Phosphorus and Vitamin D!

TOTAL CONTENT:

Calcium 12,5%
Phosphorus..... 6,4%
Propylene glycol as energizer

**With
Propylene Glycol**
AS GLUCOSE INDUCER



Calcium:

- ▶ It is the most abundant macromineral in the animal's body and the one with the greatest loss at the beginning of lactation.
- ▶ It is the first need for mineral supplementation due to the blood Calcium decrease from day 1 before calving and 3 days after.
- ▶ On the first day of calving, up to 9 times more Calcium is required for the colostrum production.
- ▶ Necessary for the uterine muscle contraction during parturition and for placenta expulsion.



Phosphorus:

- ▶ Intimately related to Calcium according to appropriate ratio: 2 of Calcium per 1 of Phosphorus (2: 1).
- ▶ The decrease in Phosphorus worsens hypocalcemia.

**Calcimax[®]
Energy**

Vitamin D3:

- ▶ Essential for the proper balance of Calcium and Phosphorus in the body.
- ▶ Promotes Calcium and Phosphorus homeostasis through intestinal absorption, mobilization, fixation and renal reabsorption pathways.

www.laboratoriosprovet.com

Provet

Eficacia ante todo.

ORAL SUSPENSION Calcimax[®] Energy






IMPROVES MILK PRODUCTION, AND THE HEALTH OF COWS AND CALVES

Calcimax[®] Energy






Designed for cow fall prevention due to milk fever (hypocalcemia).

-  Promotes the proper formation of colostrum.
-  Helps with acid-base regulation supported by Phosphorus.
-  Decreases the incidence of ketosis, mastitis, uterine prolapse, placental retention and metritis.



With Phosphorus (P) an essential ATP formation component and propylene glycol as quick source of energy.

-  Reduces negative energy balance and its effects on the cow.
-  Stimulates ovarian activity.
-  Increases the profitability of dairy farming systems.



INDICATIONS: Calcimax Energy oral suspension is indicated for the treatment of Calcium and Phosphorus deficiencies, also supports management for Vitamin D3 deficiencies and rickets.

MODE OF USE IN POSTPARTUM COW:



350 mL minutes after calving.



350 mL 12 hours after calving.



350 mL 24 hours after calving.

Each 100 mL contain: Tricalcium Phosphate: 32400 mg (Calcium ion contribution: 12500 mg; Phosphorus ion: 6460 mg). Vitamin D3: 10,000 I.U. Vitamin A Palmitate: 51000 I.U. Excipients q.s.100 mL (includes Propylene Glycol)

PRESENTATION:
350 mL bottle



Eficacia ante todo.