

ANTICOCCIDIAL GRANULATED POWDER

Coccigan[®]

20% Amprolium

COMPOSITION:

Each 100 g of Coccigan[®] contain:
Amprolium.....20 g.

Parallel therapy which allows a complete treatment of internal parasites

When deworming your animals, be aware that coccidias can produce as much or even more damage to your animals than others parasites.

Clinical signs: They might go unnoticed or be confused with other gastrointestinal and pulmonary parasites. In acute disease bloody diarrhea can be observed.

Effective action against:

Avian coccidiosis

Eimeria tenella
Eimeria acervulina
Eimeria mivati
Eimeria máxima
Eimeria brunetti

Bovine coccidiosis

Eimeria bovis
Eimeria zumii



Product presentation:

Dispensing box for fifteen 25 gram envelopes.

Dose:

•**Poultry:** Oral route, previous dissolution on drinking water. Administer: 1 gram of product in 1 liter of drinking water daily for 7 consecutive days. Then, 1 gram of product in 4 liters of drinking water for the next 7 consecutive days.

Medicated water must be the only source of drink for the birds during the treatment.

Bovine and Sheep: 1 gram of product per 20 kilograms of bodyweight for 5 consecutive days. Dissolved in water or mixed with mineralized salt.



Efficiency above all.

Coccigan[®]

ANTICOCCIDIAL GRANULATED POWDER

Can be applied in salt

CURATIVE TREATMENT WITH Coccigan[®] - MIXED WITH MINERALIZED SALT FOR BOVINE:

For the use of medication in food, it should be taken into account the mineralized salt consumption per animal and the number of animals to be treated, as follows:

Estimated daily consumption of salt	
Bodyweight (kg)	Daily consumed salt (g)
100-200	60
201-300	80
301-400	100
401-500	120

* This is an estimated consumption; it must be defined which one applies better for each scenario.

Treatment example for 60 animals of 300 kg:

Each animal daily consumes 80 g of salt. Then 60 animals consume 4800 g of salt. (80 g x 60 animals = 4800 gr of salt per day for whole population). Each animal needs 15 gr of amprolium (1g / 20 kg = 15 g / 300 kg). It means that 60 animals should consume 900 g of product (15 g x 60 animals = 900 g of Coccigan[®] per day).

In order to know how much salt is required for the preparation the amount of Coccigan[®] is subtracted from the salt amount like this:

4800 g of salt minus 900 g of Coccigan[®] is equal to 3900 g of salt plus 900 g of Coccigan[®].

It must be mixed homogeneously, from equal cores: Mix the 900 g of Coccigan[®] with 900 gr of salt until is homogeneous. Then, add 1800 g of additional salt and mix again until is homogeneous. Finally, add the remaining 1200gr of salt, until the whole mixture is completely homogenous.

COMPOSITION:

Each 100 g of Coccigan[®] contain:

Amprolium..... 20 g.

TREATMENT

CURATIVE:

Birds: 1 g/L for 7 consecutive days. 1 g/4L for the next 7 days.

Bovines and sheep: 1g/20kg for 5 consecutive days.

PREVENTIVE:

Bovines and sheep: 1 g/40kg for 10 to 20 consecutive days.

PRODUCT PRESENTATION:

Dispensing box for fifteen 25 gram envelopes.



Provet

Efficiency above all.