


Oral emulsion **Flativet**®

Composition :

Each 100ml of **Flativet**® contains
Dimethylpolysiloxane 30 g

Defoamer and antiflatulent

FLATIVET, by reducing the surface tension of bubbles inside the Rumen, **FLATIVET** inhibits them from forming again and liberates the leftover gas, thus allowing the animal to recover rapidly.



INDICATIONS

Bloat: Mainly foam-type caused by the ingestion of legumes, excessively tender green fodder or overheated fodder.

Flativet® suspension reduces the production of gases, assuring a quick recovery.

It's the highest and safest source of fluid concentrate for the prevention and treatment of chronic and acute tympanism.

Severe gastroenteritis: Lubricates the intestinal wall; reduces friction and detachment of the mucosa through the formation of a protective layer that prevents the absorption of bacterial toxins.

Dose:

Prior dilution in 250 mL of water.

Bovines and Equines: 40 mL / 100kg weight

Sheep and Goats: 20 mL / 50 Kg weight

Dogs and Cats: 5 - 10 mL according to size

Diluted: dilute 120 mL of **Flativet**® suspension in 250 mL of water.

In practice: Dilute the **FLATIVET** bottle of 120 mL in approximately 250 mL of water, for a total dilution of 370 mL. In cattle and horses, administrate 1 mL/2.5 body weight (40 mL for each 100 Kg of weight). Administrate the leftover in the next 8 to 12 hours.



Presentation:
120 mL Bottle

Oral emulsion

Flativet®

Composition :

Each 100ml of **Flativet**® contains
Dimethylpolysiloxane 30 g

Defoamer and antifatulent

INDICATIONS:

- Prevention and control of flatulencies.
- Treatment of Timpanism.
- Helps to the emptying of the intestinal tract.
- In case of obstruction of intestinal tract by feces.

Severe gastroenteritis: Lubricates the intestinal wall; reduces friction and detachment of the mucosa through the formation of a protective layer that prevents the absorption of bacterial toxins.



BENEFITS

1. Controls the accumulation of gas in the digestive tract.
2. Prevents the provision of torsion and gastric vólvulos.
3. Acts as a lubricant of the gastro-intestinal tract.
4. Promotes the intestinal transit (no laxative effect).

Highly recommended for snoring dogs and/or breed of dogs with short nose (Brachycephalic).

Dose:

Dogs and cats: 5 - 10 mL according to size (undiluted).



Presentation:

Bottle 120 ml